

À des fins de recherche uniquement

Anticorps Recombinant de lapin anti-Phospho-TAK1 (Thr187)



Numéro de catalogue: 81785-1-RR

Informations de base

Numéro de catalogue: 81785-1-RR	Numéro d'acquisition GenBank: BC017715	Méthode de purification: Purification par protéine A
Taille: 100ul , Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI): 6885	CloneNo.: 206
Hôte: Lapin	Nom complet: mitogen-activated protein kinase kinase kinase 7	Dilutions recommandées: WB 1:20000-1:100000
Isotype: IgG	MW calculé: 579 aa, 64 kDa	
	MW observés: 75-85 kDa	

Applications

Applications testées:
WB, ELISA

Spécificité de l'espèce:
Humain

Contrôles positifs:

WB : cellules HEK-293 traitées à la calyculine A, cellules HeLa traitées à la calyculine A

Informations générales

MAP3K7 (Mitogen-activated protein kinase kinase kinase 7) is also named TAK1 and belongs to the MAP kinase kinase kinase subfamily. It plays an important role in the cascades of cellular responses evoked by changes in the environment. It has been linked to interleukin-1 receptor and tumor necrosis factor receptor signaling (PMID: 16186825). It has 4 isoforms (53-55 kDa; 64-70 kDa and 75-80 kDa) produced by alternative splicing.

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Non-treated and Calyculin A treated HEK-293 cells were subjected to SDS PAGE followed by western blot with 81785-1-RR (Phospho-TAK1 (Thr187) antibody) at dilution of 1:100000 incubated at room temperature for 1 hour. The membrane was stripped and re-blotted with Alpha Tubulin antibody as loading control.