

À des fins de recherche uniquement

Anticorps Recombinant de lapin anti-Phospho-p70(S6K) (Thr389)



Numéro de catalogue: 82373-1-RR

Informations de base

Numéro de catalogue: 82373-1-RR	Numéro d'acquisition GenBank: NM_003161	Méthode de purification: Purification par protéine A
Taille: 100ul , Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI): 6198	CloneNo.: 1G15
Hôte: Lapin	Nom complet: ribosomal protein S6 kinase, 70kDa, polypeptide 1	Dilutions recommandées: WB 1:5000-1:50000
Isotype: IgG	MW observés: 65-85 kDa	

Applications

Applications testées: WB, ELISA	Contrôles positifs: WB : cellules MCF-7 traitées à l'IGF-1,
Spécificité de l'espèce: Humain	

Informations générales

The Rps6kb1 gene encodes the 70 kDa ribosomal protein S6 kinase (p70S6K), which is a serine/threonine kinase regulated by phosphoinositide 3-kinase (PI3K)/mammalian target of rapamycin (mTOR) pathway. P70S6K plays a crucial role in controlling cell cycle, growth and survival. The PI3K/mTOR signalling pathway is one of the major mechanisms for controlling cell survival, proliferation and metabolism and is the central regulator of translation of some components of protein synthesis system. Due to alternative translation two isoform S6K1 proteins are known to exist in mammalian cells: p85 S6K1 and p70 S6K1, which is identical to p85 S6K but lacks its first 23 amino acids. In addition, mammalian cells express a second S6K1 isoform spanning 316 amino acids (p31 S6K1). mTOR is known to phosphorylate and thereby activate p70 S6K1 at T389 and p85 S6K1 at T412. (PMID: 25100792, PMID: 24970012, PMID: 21602892)

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azote de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Non-treated and IGF-1 treated MCF-7 cells were subjected to SDS PAGE followed by western blot with 82373-1-RR (Phospho-p70(S6K) (Thr389) antibody) at dilution of 1:40000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with beta actin antibody (66009-1-Ig) as loading control.