

À des fins de recherche uniquement

Anticorps Recombinant de lapin anti-TIA1



Numéro de catalogue: 82672-1-RR

Informations de base

Numéro de catalogue:	BC015944	Méthode de purification:
82672-1-RR		Purification par protéine A
Taille:	Identification du gène (NCBI):	CloneNo.:
100ul , Concentration: 1000 µg/ml by Nanodrop;	7072	3N22
Hôte:	Nom complet:	Dilutions recommandées:
Lapin	TIA1 cytototoxic granule-associated RNA binding protein	WB 1:5000-1:50000
Isotype:	MW calculé	
IgG	214 aa, 24 kDa, 43 kDa	
Immunogen Catalog Number:	MW observés:	
AG2778	38-40 kDa	

Applications

Applications testées:	Contrôles positifs:
WB, ELISA	WB : cellules Caco-2, cellules COLO 320, cellules HSC-T6, cellules Jurkat, cellules K-562, cellules MOLT-4, cellules NIH/3T3
Spécificité de l'espèce:	
Humain, rat, souris	

Informations générales

TIA1, also named as p40-TIA-1, is involved in alternative pre-RNA splicing and regulation of mRNA translation by binding to AU-rich elements (AREs) located in mRNA 3' untranslated regions (3' UTRs). It possesses nucleolytic activity against cytotoxic lymphocyte target cells. TIA1 may be involved in apoptosis. Two isoforms of this protein exist - 41kDa and 42kDa. one of these was a missense variant (P362L) in TIA1. Similar to the ALS-related disease proteins TDP-43, hnRNPA1, and FUS, TIA1 is an RNA-binding protein containing a prionlike LCD and assembles into membrane-less organelles, including SGs. Postmortem neuropathology of five TIA1 mutations carriers showed a consistent pathological signature with numerous round, hyaline, TAR DNA-binding protein 43 (TDP-43)-positive inclusions.TIA1mutations significantly increased the propensity of TIA1 protein to undergo phase transition. In live cells,TIA1mutations delayed stress granule (SG) disassembly and promoted the accumulation of non-dynamic SGs that harbored TDP-43.

Stockage


Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 82672-1-RR (TIA1 antibody) at dilution of 1:25000 incubated at room temperature for 1.5 hours.