À des fins de recherche uniquement

Anticorps Monoclonal anti-CD279 (PD-1)



Numéro de catalogue: AB-65142

Informations de base

Numéro de catalogue:

AB-65142

Taille:

100ug , 0.5 mg/ml

Hôte:

Isotype: IgG2b, kappa Numéro d'acquisition GenBank:

BC119179

Identification du gène (NCBI):

18566

Nom complet:

programmed cell death 1

Méthode de purification: Purification par protéine A

CloneNo.: RMP1-30

Excitation/Emission maxima wavelengths:

404 nm / 458 nm

Applications

Applications testées:

FC

Spécificité de l'espèce:

souris

Informations générales

Programmed cell death 1 (PD-1, also known as CD279) is an immunoinhibitory receptor that belongs to the CD28/CTLA-4 subfamily of the Ig superfamily. It is a 288 amino acid (aa) type I transmembrane protein composed of one Ig superfamily domain, a stalk, a transmembrane domain, and an intracellular domain containing an immunoreceptor tyrosine-based inhibitory motif (ITIM) as well as an immunoreceptor tyrosine-based switch motif (ITSM) (PMID: 18173375). PD-1 is expressed during thymic development and is induced in a variety of hematopoietic cells in the periphery by antigen receptor signaling and cytokines (PMID: 20636820). Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function (PMID: 19426218). It is critical for the regulation of T cell function during immunity and tolerance. Blockade of PD-1 can overcome immune resistance and also has been shown to have antitumor activity (PMID: 22658127; 23169436).


Stockage

Stockage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.09% sodium azide.

Données de validation sélectionnées



1X10^6 unstimulated or Anti-CD3 and Anti-CD28 stimulated mouse splenocytes were surface stained with 0.5 ug Atlantic Blue™ Anti-Mouse CD279 (PD-1) (AB-65142, Clone: RMP1-30). Cells were not fixed.