À des fins de recherche uniquement

## Anticorps Monoclonal anti-CD16

Numéro de catalogue: APC-65090





Informations de base	Numéro de catalogue: APC-65090	Numéro d'acquisition GenBank: BC017865	Méthode de purification: Purification par affinité				
	Taille: 100tests , 5 µl/test	Identification du gène (NCBI): 2214	CloneNo.: 3G8				
	Hôte: Mouse Isotype: Mouse IgG1, k	Nom complet: Fc fragment of IgG, low affinity IIIa, receptor (CD16a) MW calculé 254 aa, 29 kDa	Excitation/Emission maxima wavelengths: 650 nm / 660 nm				
				Applications	Applications testées: FC		
					Spécificité de l'espèce: Humain		
Informations générales	CD16 is a 50-70-kDa low affinity Fc receptor found on the surface of natural killer cells, neutrophil polymorphonuclear leukocytes, monocytes and macrophages. CD16 mediates antibody-dependent cellular cytotoxicity (ADCC) and other antibody-dependent responses, such as phagocytosis. CD16 has been identified as Fc receptors Fcy RIIIa (CD16a) and Fcy RIIIb (CD16b), encoded by two nearly identical genes, FCGR3A and the FCGR3B. Clone 3G8 recognizes both the CD16a and CD16b (PMID: 7592758).						
Stockage	Stockage: Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Tampon de stockage: PBS with 0.09% sodium azide and 0.5% BSA.						

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Données de validation sélectionnées





1X10<sup>^6</sup> human PBMCs were surface co-stained with PE Anti-Human CD3 and 5 ul APC Anti-Human CD16 (APC-65090, Clone:3G8) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.

1X10<sup>^6</sup> human PBMCs were surface co-stained with PE Anti-Human CD3 and 5 ul APC Anti-Human CD16 (APC-65090, Clone:3G8) or Mouse IgG1 Isotype Control. Cells were not fixed. CD3lymphocytes were gated.