

À des fins de recherche uniquement

Anticorps Monoclonal anti-CD279 (PD-1)



Numéro de catalogue: **APC-65119**

Informations de base

| | | |
|-----------------------------------|---|---|
| Numéro de catalogue: APC-65119 | Numéro d'acquisition GenBank: BC074740 | Méthode de purification: Purification par affinité |
| Taille: 100 tests, 5 µl/test | Identification du gène (NCBI): 5133 | CloneNo.: J110 |
| Hôte: Mouse | Nom complet: programmed cell death 1 | Excitation/Emission maxima wavelengths: 650 nm / 660 nm |
| Isotype: IgG1 | MW calculé 288 aa, 32 kDa | |

Applications

Applications testées:
FC
Spécificité de l'espèce:
Humain

Informations générales

Programmed cell death 1 (PD-1, also known as CD279) is an immunoinhibitory receptor that belongs to the CD28/CTLA-4 subfamily of the Ig superfamily. It is a 288 amino acid (aa) type I transmembrane protein composed of one Ig superfamily domain, a stalk, a transmembrane domain, and an intracellular domain containing an immunoreceptor tyrosine-based inhibitory motif (ITIM) as well as an immunoreceptor tyrosine-based switch motif (ITSM) (PMID: 18173375). PD-1 is expressed during thymic development and is induced in a variety of hematopoietic cells in the periphery by antigen receptor signaling and cytokines (PMID: 20636820). Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function (PMID: 19426218). It is critical for the regulation of T cell function during immunity and tolerance. Blockade of PD-1 can overcome immune resistance and also has been shown to have antitumor activity (PMID: 22658127; 23169436).

Stockage

Stockage:
Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.
Tampon de stockage:
Tampon à base de phosphate avec azoture de sodium à 0,09 % et gélatine à 0,1 %, pH 7,2.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1X10⁶ PHA-stimulated (5 µg/mL, overnight) human peripheral blood lymphocytes were surface stained with 5.00 µl APC Anti-Human CD279 (PD-1) (APC-65119, Clone: J110) and FITC Anti-Human CD3 (FITC-65151, Clone: UCHT1). Cells were not fixed.



1X10⁶ human peripheral blood lymphocytes were surface stained with 5.00 µl APC Anti-Human CD279 (PD-1) (APC-65119, Clone: J110) and FITC Anti-Human CD3 (FITC-65151, Clone: UCHT1). Cells were not fixed.