

À des fins de recherche uniquement

# Anticorps Monoclonal anti-CD134

Numéro de catalogue:[APC-65136](#)



## Informations de base

Numéro de catalogue:	BC065782	Méthode de purification:
APC-65136		Purification par affinité
Taille:	Identification du gène (NCBI):	CloneNo.:
100ug , 0.2 mg/ml	22163	OX-86
Hôte:	Nom complet:	Excitation/Emission maxima
Rat	tumor necrosis factor receptor	wavelengths:
Isotype:	superfamily, member 4	650 nm / 660 nm
IgG1, kappa		

## Applications

Applications testées:  
FC

Spécificité de l'espèce:  
souris

## Informations générales

CD134, also known as OX40 and TNFRSF4, is a member of the TNFR-superfamily of receptors (PMID: 2828930; 9766631). It is a type I transmembrane protein predominantly expressed on activated T cells which include CD4 and CD8 T cells, Th2, Th1, and Th17 cells, as well as regulatory T cells (Tregs) (PMID: 20307208). CD134 is activated by its cognate ligand CD134L (OX40L) and functions as a T cell co-stimulatory molecule (PMID: 26215166). CD134-CD134L interactions have been proposed as a potential therapeutic target for treating autoimmune diseases, cancer and infectious disease (PMID: 26215166; 19426222).


## Stockage

Stockage:  
Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.  
Tampon de stockage:  
Tampon à base de phosphate avec azoture de sodium à 0,09 % et gélatine à 0,1 %, pH 7,2.


For technical support and original validation data for this product please contact:  
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)      E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Données de validation sélectionnées



1X10<sup>6</sup> Con A-stimulated (3 ug/ml, 2 days) BALB/c mouse splenocytes were surface stained with Coralite®488-Conjugated Anti-Mouse CD3 (CL488-65077, Clone: 17A2) and 0.13 ug APC Anti-Mouse CD134 (APC-65136, Clone: OX-86). Cells were not fixed.



1X10<sup>6</sup> Con A-stimulated (3 ug/ml, 2 days) BALB/c mouse splenocytes were surface stained with Coralite®488-Conjugated Anti-Mouse CD3 (CL488-65077, Clone: 17A2) and 0.13 ug APC-Rat IgG1 isotype control. Cells were not fixed.