

À des fins de recherche uniquement

Anticorps Monoclonal anti-CD279 (PD-1)

Numéro de catalogue:**APC-65142** 1 Publications



Informations de base

Numéro de catalogue:	BC119179	Méthode de purification:
APC-65142		Purification par affinité
Taille:	Identification du gène (NCBI):	CloneNo.:
100ug , 0.2 mg/ml	18566	RMP1-30
Hôte:	Nom complet:	Excitation/Emission maxima
Rat	programmed cell death 1	wavelengths:
Isotype:		650 nm / 660 nm
IgG2b		

Applications

Applications testées:
FC

Demandes citées:
FC

Spécificité de l'espèce:
souris

Espèces citées:
souris

Informations générales

Programmed cell death 1 (PD-1, also known as CD279) is an immunoinhibitory receptor that belongs to the CD28/CTLA-4 subfamily of the Ig superfamily. It is a 288 amino acid (aa) type I transmembrane protein composed of one Ig superfamily domain, a stalk, a transmembrane domain, and an intracellular domain containing an immunoreceptor tyrosine-based inhibitory motif (ITIM) as well as an immunoreceptor tyrosine-based switch motif (ITSM) (PMID: 18173375). PD-1 is expressed during thymic development and is induced in a variety of hematopoietic cells in the periphery by antigen receptor signaling and cytokines (PMID: 20636820). Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function (PMID: 19426218). It is critical for the regulation of T cell function during immunity and tolerance. Blockade of PD-1 can overcome immune resistance and also has been shown to have antitumor activity (PMID: 22658127; 23169436).

Publications notables

Autrice	Pubmed ID	Journal	Application
Qing Zhang	35765095	Stem Cell Res Ther	FC

Stockage


Stockage:
Stocker à une température comprise entre 2 °C et 8 °C. Éviter toute exposition à la lumière.

Tampon de stockage:
Tampon à base de phosphate avec azoture de sodium à 0,09 % et gélatine à 0,1 %, pH 7,2.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1X10⁶ Con-A stimulated BALB/c mouse splenocytes (3 ug/ml Con-A, 3 days) were surface stained with 0.5 ug APC Anti-Mouse CD279 (PD-1) (APC-65142, Clone: RMP1-30) (blue) or isotype control antibody (black). Cells were not fixed.