

À des fins de recherche uniquement

Anticorps Monoclonal anti-CD86

Numéro de catalogue: APC-65165

1 Publications



Informations de base

Numéro de catalogue:

APC-65165

Taille:

100 tests, 5 µl/test

Hôte:

Mouse

Isotype:

IgG1, kappa

Numéro d'acquisition GenBank:

BC040261

Identification du gène (NCBI):

942

Nom complet:

CD86 molecule

MW calculé

329 aa, 38 kDa

Méthode de purification:

L'anticorps purifié est conjugué à l'allophycocyanine (APC) dans des conditions optimales. Le conjugué est purifié par chromatographie d'exclusion de taille.

CloneNo.:

BU63

Excitation/Emission maxima wavelengths:

650 nm / 660 nm

Applications

Applications testées:

FC

Demandes citées:

FC

Spécificité de l'espèce:

Humain

Espèces citées:

Humain

Informations générales

CD86 (also known as B7.2) is a costimulatory molecule belonging to the immunoglobulin superfamily. Primarily expressed on antigen-presenting cells (APCs), including B cells, dendritic cells, and macrophages, CD86 is the ligand for two proteins at the cell surface of T cells, CD28 antigen and cytotoxic T-lymphocyte-associated protein 4. Binding of CD86 with CD28 antigen is a costimulatory signal for activation of the T-cell. Binding of CD86 with cytotoxic T-lymphocyte-associated protein 4 negatively regulates T-cell activation and diminishes the immune response.

Publications notables

Autrice	Pubmed ID	Journal	Application
Xueqi Yan	35413945	Cell Death Discov	FC

Stockage

Stockage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Tampon de stockage:

PBS avec azoture de sodium à 0,1 %.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1X10⁶ human PBMCs were surface co-stained with FITC Anti-Human CD14 and 5 ul APC Anti-Human CD86 (APC-65165, Clone:BU63) or unstained. Cells were not fixed. Monocytes were gated.



1X10⁶ human PBMCs were surface stained with 5 ul APC Anti-Human CD86 (APC-65165, Clone:BU63) or unstained. Cells were not fixed. Monocytes were gated.