

À des fins de recherche uniquement

Anticorps Monoclonal anti-CD279

Numéro de catalogue:[APC-65168](#)



Informations de base

| | | |
|-----------------------|---|---|
| Numéro de catalogue: | BC074740 | Méthode de purification: |
| APC-65168 | L'anticorps purifié est conjugué à l'allophycocyanine (APC) dans des conditions optimales. Le conjugué est purifié par chromatographie d'exclusion de taille. | |
| Taille: | 5133 | CloneNo.: |
| 100tests , 10 µl/test | programmed cell death 1 | EH12.2H |
| Hôte: | MW calculé | Excitation/Emission maxima wavelengths: |
| Mouse | 288 aa, 32 kDa | 650 nm / 660 nm |
| Isotype: | | |
| IgG1, kappa | | |

Applications

Applications testées:

FC

Spécificité de l'espèce:

Humain, primates non humains

Informations générales

Programmed cell death 1 (PD-1, also known as CD279) is an immunoinhibitory receptor that belongs to the CD28/CTLA-4 subfamily of the Ig superfamily. It is a 288 amino acid (aa) type I transmembrane protein composed of one Ig superfamily domain, a stalk, a transmembrane domain, and an intracellular domain containing an immunoreceptor tyrosine-based inhibitory motif (ITIM) as well as an immunoreceptor tyrosine-based switch motif (ITSM) (PMID: 18173375). PD-1 is expressed during thymic development and is induced in a variety of hematopoietic cells in the periphery by antigen receptor signaling and cytokines (PMID: 20636820). Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function (PMID: 19426218). It is critical for the regulation of T cell function during immunity and tolerance. Blockade of PD-1 can overcome immune resistance and also has been shown to have antitumor activity (PMID: 22658127; 23169436).

Stockage

Stockage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Tampon de stockage:


PBS avec azoture de sodium à 0,1 %.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1X10⁶ human peripheral blood mononuclear cells (PBMCs) were surface stained with FITC Anti-Human CD3 (FITC-65151, clone UCHT1) and 10.00 μ l/test APC Anti-Human CD279 (APC-65168, Clone:EH12.2H7). Cells were not fixed.

1X10⁶ human peripheral blood mononuclear cells (PBMCs) were surface stained with FITC Anti-Human CD3 (FITC-65151, clone UCHT1) and APC mouse IgG1 isotype control antibody. Cells were not fixed.