

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-STK25

Numéro de catalogue:Biotin-25821 Phare



Informations de base

Numéro de catalogue:	Biotin-25821	Numéro d'acquisition GenBank:	BC015793	Méthode de purification:
Taille:	100ul , Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI):	10494	Purification par affinité contre l'antigène
Hôte:	Lapin	Nom complet:	serine/threonine kinase 25 (STE20 homolog, yeast)	Dilutions recommandées:
Isotype:	IgG	MW calculé	48 kDa	IHC 1:50-1:500
Immunogen Catalog Number:	AG22964	MW observés:	48 kDa	

Applications

Applications testées:	IHC	Contrôles positifs:
Spécificité de l'espèce:	Humain, rat, souris	IHC : tissu testiculaire de souris, tissu cérébral de souris
Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9.0; (*) À défaut, 'le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.		

Informations générales

Stockage


Stockage:
Stocker à -20 °C. Éviter toute exposition à la lumière. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées




Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using Biotin-25821 (STK25 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using Biotin-25821 (STK25 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using Biotin-25821 (STK25 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using Biotin-25821 (STK25 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using Biotin-25821 (STK25 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using Biotin-25821 (STK25 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).