À des fins de recherche uniquement

Anticorps Monoclonal anti-IL-1 β

Numéro de catalogue:CL405-65276





Informations de base	Numéro de catalogue: CL405-65276	Numéro d'acquisition GenBank: BC008678	Méthode de purification: Purification par affinité
	Taille: 100ug , 0.2 mg/mL	Identification du gène (NCBI): 3553	CloneNo.: 2A7C6
	Hôte: Mouse	Nom complet: interleukin 1, beta	Excitation/Emission maxima wavelengths: 399 nm / 422 nm
	lsotype: lgG1	MW calculé 269 aa, 31 kDa	
	Immunogen Catalog Number: HZ-1164		
Applications	Applications testées: FC		
	Spécificité de l'espèce: Humain		
Informations générales	Interleukin-1 is a pro-inflammatory cytokine with multiple biological effects. The IL-1 gene family encodes three proteins: IL-10, IL-1β and their naturally occurring inhibitor IL-1RN. Interleukin 1β (IL-1β), mainly produced by blood monocytes and tissue macrophages, has been implicated in mediating both acute and chronic inflammation. IL-1β is known to be involved in a variety of cellular activities, including cell proliferation, differentiation and apoptosis. IL-1β is emerging as a key mediator of carcinogenesis that characterizes host-environment interactions. (PMID: 8630372; 12401481; 23704929; 24618930)		
Stockage	Stockage: Store at 2-8°C. Avoid exposure to Tampon de stockage: PBS with 0.09% sodium azide ar	o light. Stable for one year after shipmen nd 0.5% BSA.	t.

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées





1X10^{^6} LPS and Brefeldin A treated human PBMCs were surface stained with CL555 Anti-Human HLA-DR then intracellularly stained with 0.2 ug CoraLite® Plus 405 Anti-Human IL-1β (CL405-65276, Clone:2A7C6), or Mouse IgG1 Isotype Control. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). Monocytes were gated.

1X10^6 LPS and Brefeldin A treated human PBMCs were intracellularly stained with 0.2 ug CoraLite® Plus 405 Anti-Human IL-1β (CL405-65276, Clone:2A7C6) (red) or Mouse IgG1 Isotype Control (blue), or unactivated and Brefeldin A treated human PBMCs were intracellularly stained with 0.2 ug CoraLite® Plus 405 Anti-Human IL-1β (CL405-65276, Clone:2A7C6) (black, dashed line). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).