

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-HMMR-Specific



Numéro de catalogue: CL488-15820

Phare

Informations de base

Numéro de catalogue:
CL488-15820

Taille:
100ul , Concentration: 1000 µg/ml by
Nanodrop;

Hôte:
Lapin

Isotype:
IgG

Numéro d'acquisition GenBank:
NM_001142556

Identification du gène (NCBI):

Nom complet:
hyaluronan-mediated motility
receptor (RHAMM)

MW calculé
84 kDa

Méthode de purification:

Purification par affinité contre
l'antigène

Excitation/Emission maxima
wavelengths:
493 nm / 522 nm

Applications

Applications testées:
FC (Intra)

Spécificité de l'espèce:
Humain, rat, souris

Informations générales

HMMR, also named RHAMM, is a hyaluronan receptor expressed in various tissues. It is involved in regulation of focal contacts and subsequent motility, thus playing roles in wound healing and tumor metastasis. Remarkably different molecular masses of RHAMM have been published. Antibodies raised against RHAMM have been reported to recognize proteins of 56, 66 and 70 kDa in the supernatant of murine 3T3 fibroblasts (PMID: 2440472). Some of the isoforms detectable by anti-RHAMM antibodies, including the 52 kDa form and a 48-49 kDa form found in 10T1/2 cells, have been speculated to be underglycosylated precursor forms of the mature RHAMM protein (PMID: 1705559).

Stockage

Stockage:

Stocker à -20 °C. Éviter toute exposition à la lumière.

Tampon de stockage:

PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1X10⁶ HepG2 cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human HMMR-Specific (CL488-15820) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).