

À des fins de recherche uniquement

# Anticorps Monoclonal anti-CD49b

Numéro de catalogue: **CL488-65065**



## Informations de base

Numéro de catalogue:

CL488-65065

Taille:

100 tests, 5 µl/test

Hôte:

Rat

Isotype:

IgM

Numéro d'acquisition GenBank:

BC065139

Identification du gène (NCBI):

16398

Nom complet:

integrin alpha 2

Méthode de purification:

Purification par affinité

CloneNo.:

DX5

Excitation/Emission maxima

wavelengths:

493 nm / 522 nm

## Applications

Applications testées:

FC

Spécificité de l'espèce:

souris

## Informations générales

Integrins are a large family of  $\alpha/\beta$  transmembrane receptors playing a key role in cell-cell interactions and cell adhesion to the extracellular matrix (PMID: 30374344). CD49b, also known as integrin alpha-2, is a 150 kDa transmembrane glycoprotein expressed on a wide variety of cell types including platelets, megakaryocytes, NK cells, NKT cells, neutrophils, and long-term activated T cells (PMID: 18802064; 2415516). Integrin alpha-2/beta-1 (also known as VLA-2) is a receptor for laminin, collagen, collagen C-propeptides, fibronectin and E-cadherin (PMID: 2557634). It is responsible for adhesion of platelets and other cells to collagens, modulation of collagen and CLG gene expression, force generation and organization of newly synthesized extracellular matrix.

## Stockage

Stockage:

Stocker à une température comprise entre 2 °C et 8 °C. Éviter toute exposition à la lumière.

Tampon de stockage:

PBS avec azoture de sodium à 0,1 % et BSA à 0,5 %, pH 7,3.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Données de validation sélectionnées



1X10<sup>6</sup> mouse splenocytes were surface stained with APC Anti-Mouse NK1.1 (CD161) (PK136) (APC-65138, Clone: PK136) and CoraLite® Plus 488-conjugated Rat IgM isotype control antibody (left) or 5.00 ul CoraLite® Plus 488-conjugated Anti-Mouse CD49b (CL488-65065, Clone: DX5) (right). Cells were not fixed.