À des fins de recherche uniquement

Anticorps Monoclonal anti-CD279 (PD-1)



Numéro de catalogue: CL488-65119

Informations de base

Numéro de catalogue:

CL488-65119

Taille:

100 tests , 5 μ l/test

Hôte:

Mouse Isotype:

Numéro d'acquisition GenBank:

BC074740

Identification du gène (NCBI):

Nom complet:

programmed cell death 1

MW calculé 288 aa, 32 kDa Méthode de purification: Purification par affinité

CloneNo.: J110

Excitation/Emission maxima wavelengths:

493 nm / 522 nm

Applications

Applications testées:

FC

Spécificité de l'espèce:

Humain

Informations générales

Programmed cell death 1 (PD-1, also known as CD279) is an immunoinhibitory receptor that belongs to the CD28/CTLA-4 subfamily of the Ig superfamily. It is a 288 amino acid (aa) type I transmembrane protein composed of one Ig superfamily domain, a stalk, a transmembrane domain, and an intracellular domain containing an immunoreceptor tyrosine-based inhibitory motif (ITIM) as well as an immunoreceptor tyrosine-based switch motif (ITSM) (PMID: 18173375). PD-1 is expressed during thymic development and is induced in a variety of hematopoietic cells in the periphery by antigen receptor signaling and cytokines (PMID: 20636820). Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function (PMID: 19426218). It is critical for the regulation of T cell function during immunity and tolerance. Blockade of PD-1 can overcome immune resistance and also has been shown to have antitumor activity (PMID: 22658127; 23169436).


Stockage

Stockage:


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS avec azoture de sodium à 0,1 % et BSA à 0,5 %, pH 7,3.

Données de validation sélectionnées



1X10^6 PHA treated human PBMCs were surface costained with CL750 Anti-Human CD3 and 5 ul CoraLite® Plus 488 Anti-Human CD279 (PD-1) (CL488-65119, Clone:J110) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.



1X10^6 PHA treated human PBMCs were surface stained with 5 ul Coralite® Plus 488 Anti-Human CD279 (PD-1) (CL488-65119, Clone:J110) (Red) or Mouse [gG1] sotype Control (blue). Unactivated cells were surface stained with CL488-65119 (black). Cells were not fixed. Lymphocytes were gated.