À des fins de recherche uniquement

Anticorps Monoclonal anti-CD154

Numéro de catalogue:CL488-65129




Informations de base	Numéro de catalogue: CL488-65129	Numéro d'acquisition GenBank: BC 119225	Méthode de purification: Purification par affinité
	Taille: 100tests , 5 µl/test	Identification du gène (NCBI): 21947	CloneNo.: MR1
	Hôte: Armenian Hamster	Nom complet: CD40 ligand	Excitation/Emission maxima wavelengths: 493 nm / 522 nm
	Isotype: IgG		495 1111 222 1111 264
Applications	Applications testées: FC		
	Spécificité de l'espèce: souris		
Informations générales	The CD40 ligand (CD40L, TRAP, CD154), a member of the TNF superfamily of ligands, is expressed as either a 33-kd transmembrane homologue or 18-kd soluble form (sCD154). CD40L is primarily expressed on activated CD4+ T cells and on a small proportion of CD8+ T cells and platelets. It binds to CD40 on antigen-presenting cells (APC), which leads to many effects depending on the target cell type. It has been suggested that CD40/CD40L interactions regulate oxidative stress and affect various signaling pathways in both the immunological and the cardiovascular systems. The CD40/CD40L system is also involved in tumorigenesis. Its expression is tightly regulated, and abnormal levels of CD40L are associated with the pathogenesis of atheromatous plaque destabilization and thrombotic events. Multiple mutations in CD40LG gene have been identified that are associated with hyper-IgM immunodeficiency syndrome type 1.		
Stockage	Stockage: Store at 2-8°C. Avoid exposure Tampon de stockage: PBS avec azoture de sodium à	e to light. Stable for one year after shipmen 0,1 % et BSA à 0,5 %, pH 7,3.	t

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1X10^6 unstimulated or Cell Stimulation Cocktail stimulated (overnight) BALB/c mouse CD4+ T cells were surface co-stained with PE Anti-Mouse CD4 (GK1.5) and 0.5 ug Coralite® Plus 488 Anti-Mouse CD154 (CL488-65129, Clone: MR1). Cells were not fixed.