

À des fins de recherche uniquement

Anticorps Monoclonal anti-STAT2

Numéro de catalogue: CL488-66485



Informations de base

Numéro de catalogue:	Numéro d'acquisition GenBank:	Méthode de purification:
CL488-66485	BC051284	Purification par protéine G
Taille:	Identification du gène (NCBI):	CloneNo.:
100μl , Concentration: 1000 μg/ml by Nanodrop;	6773	1G12C4
Hôte:	Nom complet:	Excitation/Emission maxima
Mouse	signal transducer and activator of transcription 2, 113kDa	wavelengths:
Isotype:	MW calculé	493 nm / 522 nm
IgG1	851 aa, 98 kDa	
Immunogen Catalog Number:	MW observés:	
AG10168	113 kDa	

Applications

Applications testées:

FC (Intra)

Spécificité de l'espèce:

Humain, rat, souris

Informations générales

STAT2, also named as p113, belongs to the transcription factor STAT family. It is a signal transducer and activator of transcription that mediates signaling by type I IFNs (IFN-alpha and IFN-beta). Following type I IFN binding to cell surface receptors, Jak kinases (TYK2 and JAK1) are activated, leading to tyrosine phosphorylation of STAT1 and STAT2. The phosphorylated STATs dimerize and associate with ISGF3G/IRF-9 to form a complex termed ISGF3 transcription factor, that enters the nucleus. ISGF3 binds to the IFN stimulated response element (ISRE) to activate the transcription of IFN stimulated genes, which drive the cell in an antiviral state. It also interacts with CRSP2, CRSP6, Simian virus 5 protein V, rabies virus phosphoprotein, IFNAR1 and IFNAR2. Its interaction with dengue virus NS5 inhibits the phosphorylation of STAT2, and, when all viral proteins are present (polyprotein), STAT2 is targeted for degradation. The calculated molecular weight of STAT2 is 98 kDa, but phosphorylated STAT2 is about 100-113 kDa.

Stockage

Stockage:

Stocker à -20 °C. Éviter toute exposition à la lumière. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.


L'aliquotage n'est pas nécessaire pour le stockage à -20°C

*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1X10⁶ Ramos cells were intracellularly stained with 0.4 ug Coralite® Plus 488 Anti-Human STAT2 (CL488-66485, Clone:1G12C4) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).