

À des fins de recherche uniquement

Anticorps Monoclonal anti-CD40L/CD154

Numéro de catalogue:[CL488-66502](#)



Informations de base

Numéro de catalogue:	BC071754	Méthode de purification:
CL488-66502	Purification par protéine A	
Taille:	Identification du gène (NCBI):	CloneNo.:
100ul , Concentration: 1000 µg/ml by Nanodrop;	959	1E6D10
Hôte:	Nom complet:	Excitation/Emission maxima
Mouse	CD40 ligand	wavelengths:
Isotype:	MW calculé	493 nm / 522 nm
IgG2a	261 aa, 29 kDa	
Immunogen Catalog Number:	MW observés:	
AG24941	32 kDa	

Applications

Applications testées:

FC (Intra)

Spécificité de l'espèce:

Humain

Informations générales

The CD40 ligand (CD40L, TRAP, CD154), a member of the TNF superfamily of ligands, is expressed as either a 33-kd transmembrane homologue or 18-kd soluble form (sCD154). CD40L is primarily expressed on activated CD4+ T cells and on a small proportion of CD8+ T cells and platelets. It binds to CD40 on antigen-presenting cells (APC), which leads to many effects depending on the target cell type. It has been suggested that CD40/CD40L interactions regulate oxidative stress and affect various signaling pathways in both the immunological and the cardiovascular systems. The CD40/CD40L system is also involved in tumorigenesis. Its expression is tightly regulated, and abnormal levels of CD40L are associated with the pathogenesis of atherosomatous plaque destabilization and thrombotic events. Multiple mutations in CD40LG gene have been identified that are associated with hyper-IgM immunodeficiency syndrome type 1.

Stockage

Stockage:

Stocker à -20 °C. Éviter toute exposition à la lumière. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.


L'aliquotage n'est pas nécessaire pour le stockage à -20°C

*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1X10⁶ Jurkat cells were intracellularly stained with 0.4 ug Coralite® Plus 488 Anti-Human CD40L/CD154 (CL488-66502, Clone:1E6D10) (red) or 0.4 ug isotype control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).