

À des fins de recherche uniquement

Anticorps Monoclonal anti-HOXA7

Numéro de catalogue: CL488-67112



Informations de base

Numéro de catalogue: CL488-67112	Numéro d'acquisition GenBank: BC148692	Méthode de purification: Purification par protéine G
Taille: 100ul, Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI): 3204	CloneNo.: 2A7H6
Hôte: Mouse	Nom complet: homeobox A7	Dilutions recommandées: IF 1:50-1:500
Isotype: IgG1	MW calculé: 230 aa, 25 kDa	Excitation/Emission maxima wavelengths: 493 nm / 522 nm
Immunogen Catalog Number: AG19025	MW observés: 20 kDa	

Applications

Applications testées: IF	Contrôles positifs: IF : cellules HepG2,
Spécificité de l'espèce: Humain	

Informations générales

HOX genes play a fundamental role in the development of the vertebrate central nervous system, heart, axial skeleton, limbs, gut, urogenital tract and external genitalia. The homeobox gene Hoxa-1 is transcriptionally regulated by retinoic acid (RA) and encodes a transcription factor, which has been shown to play important roles in cell differentiation and embryogenesis. Hoxa-1 is also expressed in cancers, such as mammary tumors, though it is not expressed in normal gland or in precancerous mammary tissues. At embryonic stages, Hoxa-2 is expressed in the mesenchyme and epithelial cells of palate, however its expression is restricted to the tips of the growing palatal shelves. Hoxa-2 protein is predominantly expressed in the nuclei of cells in the ventral mantle region of the developing embryo. In the developing and adult mouse spinal cord, Hoxa-2 protein may contribute to dorsal-ventral patterning and/or to the specification of neuronal phenotype. Hoxa-7 functions as a potent transcriptional repressor and its action as such requires several domains, including both activator and repressor regions. Hoxa-7 is expressed in the fetal liver, lung, skeletal muscle, kidney, pancreas and placenta.

Stockage

Stockage:
Stocker à -20 °C. Éviter toute exposition à la lumière.
Tampon de stockage:
PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.
L'aliquotage n'est pas nécessaire pour le stockage à -20C

***** Les 20ul contiennent 0,1% de BSA.**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite® Plus 488 HOXA7 antibody (CL488-67112, Clone: 2A7H6) at dilution of 1:200.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite® Plus 488 HOXA7 antibody (CL488-67112, Clone: 2A7H6) at dilution of 1:200.