

À des fins de recherche uniquement

Anticorps Monoclonal anti-PSME1

Numéro de catalogue:**CL488-67928**



Informations de base

Numéro de catalogue:	BC007503	Méthode de purification:
CL488-67928		Purification par protéine G
Taille:	Identification du gène (NCBI):	CloneNo.:
100ul , Concentration: 1000 µg/ml by Nanodrop;	5720	1F7E2
Hôte:	Nom complet:	Excitation/Emission maxima
Mouse	proteasome (prosome, macropain) activator subunit 1 (PA28 alpha)	wavelengths:
Isotype:	MW calculé	493 nm / 522 nm
IgG1	29 kDa	
Immunogen Catalog Number:	MW observés:	
AG30549	29 kDa	

Applications

Applications testées:

FC (Intra)

Spécificité de l'espèce:

Humain, souris

Informations générales

The principal function of the proteasome is targeted degradation of intracellular proteins. Activity of the 20S proteasome is controlled by regulatory complexes that bind to the ends of the cylindrical proteasome. 11S regulator (REG or PA28), is a complex of 28 kDa subunits that is thought to activate proteasomes toward the production of antigenic peptides. Human PSME1 and PSME2 genes encode the two proteasome activators PA28 alpha and beta, respectively, which have been implicated in antigen processing for loading class I MHC molecules. The PA28 activator complex enhances the generation of class I binding peptides by altering the cleavage pattern of the proteasome.

Stockage

Stockage:

Stocker à -20 °C. Éviter toute exposition à la lumière. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.


L'aliquotage n'est pas nécessaire pour le stockage à -20°C

*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1×10^6 HepG2 cells were intracellularly stained with 0.4 ug CoralLite® Plus 488 Anti-Human PSME1 (CL488-67928, Clone:1F7E2) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).