

À des fins de recherche uniquement

Anticorps Monoclonal anti-STK11/LKB1



Numéro de catalogue: **CL488-68016**

Informations de base

Numéro de catalogue: CL488-68016	Numéro d'acquisition GenBank: BC007981	Méthode de purification: Purification par protéine G
Taille: 100ul , Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI): 6794	CloneNo.: 2D11E5
Hôte: Mouse	Nom complet: serine/threonine kinase 11	Dilutions recommandées: IF 1:50-1:500
Isotype: IgG1	MW calculé 49 kDa	Excitation/Emission maxima wavelengths: 493 nm / 522 nm
	MW observés: 50-55 kDa	

Applications

Applications testées: FC (Intra), IF	Contrôles positifs: IF : cellules NIH/3T3,
Spécificité de l'espèce: Humain, rat, souris	

Informations générales

STK11 (serine/threonine-protein kinase 11) is also named as LKB1, PJS, and belongs to the protein kinase superfamily. It controls the activity of AMP-activated protein kinase (AMPK) family members, thereby playing a role in various processes such as cell metabolism, cell polarity, apoptosis and DNA damage response. The tumour suppressor protein LKB1 is a serine/threonine kinase that has been causally linked to Peutz-Jeghers syndrome (PJS). Defects in STK11 are a cause of Peutz-Jeghers syndrome (PJS) and defects in STK11 have been associated with testicular germ cell tumor (TGCT) and some sporadic cancers, especially lung cancers. STK11 has 2 isoforms with MW of 49 kDa and 45 kDa, and can be detected as 50-54 kDa after posttranslational modification.

Stockage

Stockage:
Stocker à -20 °C. Éviter toute exposition à la lumière. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.
L'aliquotage n'est pas nécessaire pour le stockage à -20C

***** Les 20ul contiennent 0,1% de BSA.**



For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1×10^6 HeLa cells were intracellularly stained with 0.4 μg CoraLite® Plus 488 Anti-Human STK11/LKB1 (CL488-68016, Clone:2D11E5) (red), or 0.4 μg x. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).

Immunofluorescent analysis of (4% PFA) fixed NIH/3T3 cells using CoraLite® Plus 488 STK11 antibody (CL488-68016, Clone: 2D11E5) at dilution of 1:200.