

À des fins de recherche uniquement

Anticorps Monoclonal anti-INO80

Numéro de catalogue: **CL488-68064**



Informations de base

| | | |
|---|---|---|
| Numéro de catalogue: CL488-68064 | Numéro d'acquisition GenBank: BC146785 | Méthode de purification: Purification par protéine G |
| Taille: 100ul, Concentration: 1000 µg/ml by Nanodrop; | Identification du gène (NCBI): 54617 | CloneNo.: 1F11C5 |
| Hôte: Mouse | Nom complet: INO80 homolog (S. cerevisiae) | Excitation/Emission maxima wavelengths: 493 nm / 522 nm |
| Isotype: IgG1 | MW calculé: 1556 aa, 177 kDa | |
| Immunogen Catalog Number: AG20265 | MW observés: 177 kDa | |

Applications

Applications testées:
FC (Intra)

Spécificité de l'espèce:
Humain, rat, souris

Informations générales

DNA helicase INO80, also named as INO80 or hINO80, is a 1,556 amino acid protein, which contains one helicase C-terminal domain, one helicase ATP-binding domain, and one DBINO domain. INO80 co-localizes with PCMA at replication forks during S-phase in the nucleus and belongs to the SNF2 helicase family. INO80 is a component of the chromatin remodeling INO80 complex and is widely expressed in many tissues, such as brain, liver and pancreas. INO80 is involved in transcription regulation.

Stockage

Stockage:
Stocker à -20 °C. Éviter toute exposition à la lumière. Stable pendant un an après l'expédition.

Tampon de stockage:
PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.
L'aliquotage n'est pas nécessaire pour le stockage à -20C

***** Les 20ul contiennent 0,1% de BSA.**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1X10⁶ PC-12 cells were intracellularly stained with 0.8 ug CoraLite® Plus 488 Anti-Human INO80 (CL488-68064, Clone:1F11C5) (red), or 0.8 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).