À des fins de recherche uniquement

Anticorps Monoclonal anti-CD134

Numéro de catalogue: CL555-65136



Informations de base

Numéro de catalogue:

CL555-65136

Taille:

100ug , 0.5 mg/ml

Hôte:

Isotype: IgG1, kappa Numéro d'acquisition GenBank:

BC065782

Identification du gène (NCBI):

22163

Nom complet:

tumor necrosis factor receptor superfamily, member 4

Méthode de purification: Purification par affinité

CloneNo.: OX-86

Excitation/Emission maxima

wavelengths: 554 nm / 570 nm

Applications

Applications testées:

FC

Spécificité de l'espèce:

souris

Informations générales

CD134, also known as OX40 and TNFRSF4, is a member of the TNFR-superfamily of receptors (PMID: 2828930; 9766631). It is a type I transmembrane protein predominantly expressed on activated T cells which include CD4 and CD8 T cells, Th2, Th1, and Th17 cells, as well as regulatory T cells (Tregs) (PMID: 20307208). CD134 is activated by its cognate ligand CD134L (OX40L) and functions as a T cell co-stimulatory molecule (PMID: 26215166). CD134-CD134L interactions have been proposed as a potential therapeutic target for treating autoimmune diseases, cancer and infectious disease (PMID: 26215166; 19426222).

Stockage


Stockage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Tampon de stockage:

PBS with 0.09% sodium azide.

Données de validation sélectionnées



1X10^6 unstimulated or anti-CD3/CD28-stimulated mouse splenocytes were surface co-stained with CoraLite® Plus 647 Anti-Mouse CD3 (17A2) and 0.5 ug CoraLite® Plus 555 Anti-Mouse CD134 (CL555-65136, Clone:OX-86). Cells were not fixed.