À des fins de recherche uniquement

Anticorps Monoclonal anti-CD8

Numéro de catalogue: CL555-65204



Informations de base

Numéro de catalogue:

CL555-65204

Taille:

100tests , 5 ul/test

Hôte: Mouse Isotype: lgG2a

Numéro d'acquisition GenBank:

BC025715

Identification du gène (NCBI):

Nom complet: CD8a molecule MW calculé 235 aa, 26 kDa

Méthode de purification: Purification par protéine A

CloneNo.: UCHT4

Excitation/Emission maxima wavelengths: 554 nm / 570 nm

Applications

Applications testées:

FC

Spécificité de l'espèce:

Humain

Informations générales

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8a or as a heterodimer of CD8a and CD8β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8αβ heterodimer while CD8αα homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR) $\gamma\delta$ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).

Stockage


Stockage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Tampon de stockage:

PBS with 0.09% sodium azide and 0.5% BSA.

Données de validation sélectionnées



1X10^6 human PBMCs were surface co-stained with APC Anti-Human CD3 and 5 ul Coralite® Plus 555 Anti-Human CD8 (CL555-65204, Clone:UCHT4) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul CoraLite® Plus 555 Anti-Human CD8 (CL555-65204, Clone:UCHT4) (red) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.