

À des fins de recherche uniquement

# Anticorps Monoclonal anti-Cytokeratin 18

Numéro de catalogue:**CL555-66187**



## Informations de base

Numéro de catalogue:	BC000180	Méthode de purification:
CL555-66187		Purification par protéine A
Taille:	Identification du gène (NCBI):	CloneNo.:
100ul , Concentration: 1000 µg/ml by Nanodrop;	3875	1G11C4
Hôte:	Nom complet:	Dilutions recommandées:
Mouse	keratin 18	IF 1:50-1:500
Isotype:	MW calculé	Excitation/Emission maxima
IgG1	48 kDa	wavelengths:
Immunogen Catalog Number:		557 nm / 570nm
AG1260		

## Applications

Applications testées:	Contrôles positifs:
FC (Intra), IF	IF : cellules HepG2,
Spécificité de l'espèce:	
Humain	

## Informations générales

Keratins are a large family of proteins that form the intermediate filament cytoskeleton of epithelial cells, which are classified into two major sequence types. Type I keratins are a group of acidic intermediate filament proteins, including K9-K23, and the hair keratins Ha1-Ha8. Type II keratins are the basic or neutral counterparts to the acidic type I keratins, including K1-K8, and the hair keratins, Hb1-Hb6. KRT18, also named as CYK18, PIG46 and K18, is the most commonly found members of the intermediate filament gene family. KRT18 is a type I keratin which is found primarily in non squamous epithelia and is present in a majority of adenocarcinomas and ductal carcinomas but not in squamous cell carcinomas.

## Stockage


**Stockage:**  
Stocker à -20 °C. Éviter toute exposition à la lumière.  
**Tampon de stockage:**  
PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.  
**L'aliquotage n'est pas nécessaire pour le stockage à -20°C**

\*\*\* Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:  
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)      E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Données de validation sélectionnées



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CoraLite®555 Cytokeratin 18 antibody (CL555-66187, Clone: 1G11C4) at dilution of 1:200.



1X10<sup>6</sup> HeLa cells were intracellularly stained with 0.2 ug CoraLite®555 Anti-Human Cytokeratin 18 (CL555-66187, Clone:1G11C4) (red), or 0.2 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).