

À des fins de recherche uniquement

# Anticorps Monoclonal anti-SUMO2/3

Numéro de catalogue: **CL555-67154** **Phare**



## Informations de base

<b>Numéro de catalogue:</b> CL555-67154	<b>Numéro d'acquisition GenBank:</b> BC008450	<b>Méthode de purification:</b> Purification par protéine A
<b>Taille:</b> 100ul , Concentration: 1000 µg/ml by Nanodrop;	<b>Identification du gène (NCBI):</b> 6613	<b>CloneNo.:</b> 1A1B3
<b>Hôte:</b> Mouse	<b>Nom complet:</b> SMT3 suppressor of mif two 3 homolog 2 (S. cerevisiae)	<b>Excitation/Emission maxima wavelengths:</b> 557 nm / 570nm
<b>Isotype:</b> IgG2b	<b>MW calculé</b> 11 kDa	
<b>Immunogen Catalog Number:</b> AG28672	<b>MW observés:</b> 18 kDa	

## Applications

**Applications testées:**  
FC (Intra)

**Spécificité de l'espèce:**  
Humain, rat, souris

## Informations générales

Ubiquitin is most famous for its function in targeting proteins for degradation by the 26S proteasome, ubiquitin needs to be attached to a substrate in chains (polyubiquitylation) before being recognized by proteasome. Similarly, SUMO (small ubiquitin-related modifier) can be linked to substrates in chains (polysumoylation), SUMO modification has been implicated in many important cellular processes including the control of genome stability, signal transduction, targeting to and formation of nuclear compartments, cell cycle and meiosis. There are 4 confirmed SUMO isoforms in human, SUMO-1, SUMO-2, SUMO-3 and SUMO-4. SUMO-2 and SUMO-3 are nearly identical but are distinct from SUMO-1. SUMO2/3 conjugation was recently widely involved in neuroprotective activities. A substitution (M55V) of SUMO4 was strongly associated with the pathogenesis of type 1 diabetes (T1D) involving NF kappa B related mechanisms.

## Stockage

**Stockage:**  
Stocker à -20 °C. Éviter toute exposition à la lumière. Stable pendant un an après l'expédition.


**Tampon de stockage:**  
PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.  
L'aliquotage n'est pas nécessaire pour le stockage à -20C

**\*\*\* Les 20ul contiennent 0,1% de BSA.**

For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)      E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Données de validation sélectionnées



1X10<sup>6</sup> Jurkat cells were intracellularly stained with 0.4 ug CoraLite®555 Anti-Human SUMO2/3 (CL555-67154, Clone:1A1B3) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).