

À des fins de recherche uniquement

Anticorps Recombinant de lapin anti- Phospho-Hamartin/TSC1 (Ser511)



Numéro de catalogue:**CL555-80340**

Informations de base

Numéro de catalogue:	Numéro d'acquisition GenBank:	Méthode de purification:
CL555-80340	NM_000368	Purification par protéine A
Taille:	Identification du gène (NCBI):	CloneNo.:
100ul , Concentration: 1000 µg/ml by Nanodrop;	7248	3N6
Hôte:	Nom complet:	Excitation/Emission maxima wavelengths:
Lapin	tuberous sclerosis 1	557 nm / 570nm
Isotype:	MW calculé	
IgG	130 kDa	
	MW observés:	
	150 kDa	

Applications

Applications testées:
FC (Intra)

Spécificité de l'espèce:
Humain, souris

Informations générales

TSC1, also named as Hamartin, is a tumor suppressor gene syndrome whose manifestations can include seizures, mental retardation, autism, and tumors in the brain, retina, kidney, heart, and skin. The tuberous sclerosis 1 (TSC1)/TSC2 tumor suppressor complex serves as a repressor of the mTOR pathway, and disruption of TSC1/TSC2 complex function may contribute to tumorigenesis. TSC1 is phosphorylated by IKK β at Ser487 and Ser511 in vitro and in vivo, and phosphorylation at these sites enhances dissociation of the TSC1/TSC2 complex, which in turn induces mTOR activation. (PMID: 17693255). TSC1 is also threonine-phosphorylated at three sites (Thr417, Ser584, and Thr1047) in a reaction catalyzed by cyclin-dependent kinase 1 (CDK1) (PMID: 14551205).

Stockage

Stockage:
Stocker à -20 °C. Éviter toute exposition à la lumière. Stable pendant un an après l'expédition.


Tampon de stockage:
PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.
L'aliquotage n'est pas nécessaire pour le stockage à -20°C

*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1X10⁶ HeLa cells untreated (dashed lines) or treated with Calyculin A (red) were intracellularly stained with 0.5 ug Coralite®555 Anti-Human Phospho-Hamartin/TSC1 (Ser511) (CL555-80340, Clone:3N6) (red), or 0.5 ug Control Antibody (blue). Cells were fixed with 4% PFA and permeabilized with 50% MeOH.

1X10⁶ HeLa cells untreated (left) or treated with Calyculin A (right) were intracellularly stained with 0.5 ug Coralite®555 Anti-Human Phospho-Hamartin/TSC1 (Ser511) (CL555-80340, Clone:3N6) (red), or 0.5 ug Control Antibody (blue). Cells were fixed with 4% PFA and permeabilized with 50% MeOH.