

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-Beta Tubulin



Numéro de catalogue:**CL594-10068**

Informations de base

Numéro de catalogue:	BC000748	Méthode de purification:
CL594-10068		Purification par affinité contre l'antigène
Taille:	Identification du gène (NCBI):	Dilutions recommandées:
100ul , Concentration: 1000 µg/ml by Nanodrop;	10381	IF 1:50-1:500
Hôte:	Nom complet:	Excitation/Emission maxima wavelengths:
Lapin	tubulin, beta 3	588 nm / 604 nm
Isotype:	MW calculé	
IgG	450 aa, 50 kDa	
Immunogen Catalog Number:	MW observés:	
AG0117	50-55 kDa	

Applications

Applications testées:	Contrôles positifs:
FC, IF	IF : cellules HeLa,
Spécificité de l'espèce:	
Humain, rat, souris	

Informations générales

There are five tubulins in human cells: alpha, beta, gamma, delta, and epsilon. Tubulins are conserved across species. They form heterodimers, which multimerize to form a microtubule filament. An alpha and beta tubulin heterodimer is the basic structural unit of microtubules. The heterodimer does not come apart, once formed. The alpha and beta tubulins, which are each about 55 kDa MW, are homologous but not identical. Alpha, beta, and gamma tubulins have all been used as loading controls. Tubulin expression may vary according to resistance to antimicrobial and antimitotic drugs.

Stockage


Stockage:
Stocker à -20 °C. Éviter toute exposition à la lumière. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.
L'aliquotage n'est pas nécessaire pour le stockage à -20°C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



WWW.PTGLAB.COM



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using Coralite®594 Beta Tubulin antibody (CL594-10068) at dilution of 1:200.

1X10⁶ HeLa cells were intracellularly stained with 0.4 ug Coralite®594 Anti-Human Beta Tubulin (CL594-10068) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).