

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-NF- κ B p65



Numéro de catalogue: **CL594-10745**

Phare

2 Publications

Informations de base

Numéro de catalogue: CL594-10745	Numéro d'acquisition GenBank: BC011603	Méthode de purification: Purification par affinité contre l'antigène
Taille: 100ul , Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI): 5970	Dilutions recommandées: IF 1:50-1:500
Hôte: Lapin	Nom complet: v-rel reticuloendotheliosis viral oncogene homolog A (avian)	Excitation/Emission maxima wavelengths: 588 nm / 604 nm
Isotype: IgG	MW calculé: 65 kDa	
Immunogen Catalog Number: AG1199	MW observés: 65 kDa	

Applications

Applications testées: IF	Contrôles positifs: IF : cellules HepG2,
Demandes citées: IF	
Spécificité de l'espèce: Humain, porc, rat, souris	
Espèces citées: souris	

Informations générales

Nuclear factor κ B (NF- κ B) is a sequence-specific DNA-binding protein complex which regulates the expression of viral genomes, including the human immunodeficiency virus, and a variety of cellular genes, particularly those involved in immune and inflammatory responses. The members of the NF- κ B family in mammalian cells include the proto-oncogene c-Rel, p50/p105 (NF κ B1), p65 (RelA), p52/p100 (NF κ B2), and RelB. All of these proteins share a conserved 300-amino acid region known as the Rel homology domain which is responsible for DNA binding, dimerization, and nuclear translocation of NF- κ B. The p65 subunit is a major component of NF- κ B complexes and is responsible for trans-activation. NF- κ B heterodimeric p65-p50 and p65-c-Rel complexes are transcriptional activators. The NF- κ B p65-p65 complex appears to be involved in invasion-mediated activation of IL-8 expression. The inhibitory effect of I κ B upon NF- κ B the cytoplasm is exerted primarily through the interaction with p65. p65 shows a weak DNA-binding site which could contribute directly to DNA binding in the NF- κ B complex. It associates with chromatin at the NF- κ B promoter region via association with DDX1. This antibody is a rabbit polyclonal antibody raised against residues near the N terminus of human RELA.

Publications notables

Autrice	Pubmed ID	Journal	Application
Zhiyuan Yu	36091713	Front Med (Lausanne)	IF
Zhiyuan Yu	37036417	Invest Ophthalmol Vis Sci	IF

Stockage


Stockage:
Stocker à -20 °C. Éviter toute exposition à la lumière.
Tampon de stockage:
PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.
L'aliquotage n'est pas nécessaire pour le stockage à -20C

***** Les 20ul contiennent 0,1% de BSA.**

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using CoraLite®594 NF-κB p65 antibody (CL594-10745) at dilution of 1:100, CoraLite®488 Beta Actin antibody (CL488-66009, Clone: 2D4H5, green), DAPI (Blue).