

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-Cytokeratin 18

Numéro de catalogue:**CL594-10830**



Informations de base

Numéro de catalogue:	BC000180	Méthode de purification:
CL594-10830		Purification par affinité contre l'antigène
Taille:	Identification du gène (NCBI):	Dilutions recommandées:
100ul , Concentration: 1000 µg/ml by Nanodrop;	3875	WB 1:1000-1:4000 IF 1:50-1:500
Hôte:	Nom complet:	Excitation/Emission maxima wavelengths:
Lapin	keratin 18	588 nm / 604 nm
Isotype:	MW calculé	
IgG	48 kDa	
Immunogen Catalog Number:	MW observés:	
AG1260	48 kDa	

Applications

Applications testées:	Contrôles positifs:
FC (Intra), IF, WB	WB : cellules A431, cellules HeLa, cellules K-562, cellules MCF-7
Spécificité de l'espèce:	
Humain, souris	IF : cellules HepG2, cellules HeLa

Informations générales

Keratins are a large family of proteins that form the intermediate filament cytoskeleton of epithelial cells, which are classified into two major sequence types. Type I keratins are a group of acidic intermediate filament proteins, including K9-K23, and the hair keratins Ha1-Ha8. Type II keratins are the basic or neutral counterparts to the acidic type I keratins, including K1-K8, and the hair keratins, Hb1-Hb6. KRT18, also named as CYK18, PIG46 and K18, is the most commonly found members of the intermediate filament gene family. KRT18 is a type I keratin which is found primarily in non squamous epithelia and is present in a majority of adenocarcinomas and ductal carcinomas but not in squamous cell carcinomas.

Stockage


Stockage:
Stocker à -20°C. Éviter toute exposition à la lumière.
Tampon de stockage:
PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with CL594-10830 (Cytokeratin 18 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using Coralite®594 Cytokeratin 18 antibody (CL594-10830) at dilution of 1:200.



1X10⁶ HeLa cells were intracellularly stained with 0.4 ug Coralite®594 Anti-Human Cytokeratin 18 (CL594-10830) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).