

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-TIA1



Numéro de catalogue: **CL594-12133**

Phare

Informations de base

Numéro de catalogue: CL594-12133	Numéro d'acquisition GenBank: BC015944	Méthode de purification: Purification par affinité contre l'antigène
Taille: 100ul , Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI): 7072	Dilutions recommandées: IF 1:50-1:500
Hôte: Lapin	Nom complet: TIA1 cytotoxic granule-associated RNA binding protein	Excitation/Emission maximum wavelengths: 588 nm / 604 nm
Isotype: IgG	MW calculé 214 aa, 24 kDa, 43 kDa	
Immunogen Catalog Number: AG2778	MW observés: 40 kDa	

Applications

Applications testées: IF	Contrôles positifs: IF : cellules LO2,
Spécificité de l'espèce: Humain, rat, souris	

Informations générales

TIA1, also named as p40-TIA-1, is involved in alternative pre-RNA splicing and regulation of mRNA translation by binding to AU-rich elements (AREs) located in mRNA 3' untranslated regions (3' UTRs). It possesses nucleolytic activity against cytotoxic lymphocyte target cells. TIA1 may be involved in apoptosis. Two isoforms of this protein exist - 41kDa and 42kDa. one of these was a missense variant (P362L) in TIA1. Similar to the ALS-related disease proteins TDP-43, hnRNPA 1, and FUS, TIA1 is an RNA-binding protein containing a prionlike LCD and assembles into membrane-less organelles, including SGs. Postmortem neuropathology of five TIA1 mutations carriers showed a consistent pathological signature with numerous round, hyaline, TAR DNA-binding protein 43 (TDP-43)-positive inclusions. TIA1 mutations significantly increased the propensity of TIA1 protein to undergo phase transition. In live cells, TIA1 mutations delayed stress granule (SG) disassembly and promoted the accumulation of non-dynamic SGs that harbored TDP-43. Moreover, TDP-43 in SGs became less mobile and insoluble.

Stockage

Stockage:
Stocker à -20 °C. Éviter toute exposition à la lumière.
Tampon de stockage:
PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Immunofluorescent analysis of (4% PFA) fixed L02 cells using Coralite® 594 TIA1 antibody (CL594-12133) at dilution of 1:200.