

À des fins de recherche uniquement

Anticorps Monoclonal anti-BRD8

Numéro de catalogue: **CL594-60121** **Phare**



Informations de base

Numéro de catalogue: CL594-60121	Numéro d'acquisition GenBank: BC008039	Méthode de purification: Purification par protéine G
Taille: 100ul , Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI): 10902	CloneNo.: 5B6D6
Hôte: Mouse	Nom complet: bromodomain containing 8	Dilutions recommandées: IF 1:50-1:500
Isotype: IgG1	MW calculé 120 kDa	Excitation/Emission maxima wavelengths: 588 nm / 604 nm
Immunogen Catalog Number: AG0790	MW observés: 120 kDa	

Applications

Applications testées: IF	Contrôles positifs: IF : cellules HepG2,
Spécificité de l'espèce: Humain	

Informations générales

BRD8 is one of the bromodomain-containing proteins, which is an acetylated lysine-binding domain and thought to be involved in regulation of protein acetylation and/or HAT (histone acetyl transferase) activity [PMID: 12963728]. It is also reported to be an accessory subunit of NuA4 HAT complex (also known as TRRAP/TIP60 complex) through biochemical purification [PMID: 16049979]. In addition, it can bind to thyroid hormone receptor-β and/or retinoid X receptor, and can act as a coactivator of the nuclear hormone receptor-mediated transcription in reporter assays [PMID: 10517671].

Stockage


Stockage:
Stocker à -20 °C. Éviter toute exposition à la lumière. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.
L'aliquotage n'est pas nécessaire pour le stockage à -20C

***** Les 20ul contiennent 0,1% de BSA.**

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CL594-60121 (BRD8 antibody) at dilution of 1:100.