

À des fins de recherche uniquement

Anticorps Monoclonal anti-AK2

Numéro de catalogue:**CL594-66127**



Informations de base

Numéro de catalogue:	Numéro d'acquisition GenBank:	Méthode de purification:
CL594-66127	BC009405	Purification par protéine A
Taille:	Identification du gène (NCBI):	CloneNo.:
100ul , Concentration: 1000 µg/ml by 204		3H2C10
Nanodrop;	Nom complet:	Dilutions recommandées:
	adenylate kinase 2	IF 1:50-1:500
Hôte:	MW calculé	Excitation/Emission maxima wavelengths:
Mouse	26 kDa	588 nm / 604 nm
Isotype:	MW observés:	
IgG2a	29 kDa	
Immunogen Catalog Number:		
AG17911		

Applications

Applications testées:	Contrôles positifs:
FC (Intra), IF	IF : cellules HepG2,
Spécificité de l'espèce:	
Humain	

Informations générales

AK2(Adenylate kinase 2, mitochondrial) is also named as ADK2 and belongs to the adenylate kinase family. It plays a unique role in energy metabolism and energy transfer by regulating the ATP/ADP rate between the cytoplasma matrix and the mitochondria and it is increased expression during adipocyte and B cell differentiation(PMID:23020757). AK2 catalyzes the reversible transfer of a phosphoryl group from ATP to AMP(PMID:19043417). Defects in AK2 are the cause of reticular dysgenesis (RDYS)(PMID:19043416). It has 6 isoforms produced by alternative splicing.

Stockage


Stockage:
Stocker à -20 °C. Éviter toute exposition à la lumière. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CL594-66127 (AK2 antibody) at dilution of 1:100.



1X10⁶ HepG2 cells were intracellularly stained with 0.4 ug Coralite®594 Anti-Human AK2 (CL594-66127, Clone:3H2C10) (red), or 0.4 ug Mouse IgG2a Isotype Control (CL594-66360-2, Clone: K11A1B2A2) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).