

À des fins de recherche uniquement

# Anticorps Monoclonal anti-Arginase-1

Numéro de catalogue: CL594-66129

1 Publications



Antibodies | ELISA kits | Proteins  
[www.ptglab.com](http://www.ptglab.com)

## Informations de base

Numéro de catalogue:	Numéro d'acquisition GenBank:	Méthode de purification:
CL594-66129	BC005321	Purification par protéine G
Taille:	Identification du gène (NCBI):	CloneNo.:
100ul , Concentration: 1000 µg/ml by Nanodrop;	383	5D6D12
Hôte:	Nom complet:	Dilutions recommandées:
Mouse	arginase, liver	IF 1:50-1:500
Isotype:	MW calculé	Excitation/Emission maxima wavelengths:
IgG1	236aa,25 kDa; 322aa,35 kDa	588 nm / 604 nm
Immunogen Catalog Number:		
AG8810		

## Applications

Applications testées:	Contrôles positifs:
IF	IF : cellules HepG2,
Demandes citées:	
IF	
Spécificité de l'espèce:	
Humain, porc, rat, souris	
Espèces citées:	
Humain	

## Informations générales

Arginase-1 (Liver arginase) belongs to the arginase family. ARG1 is a novel immunohistochemical marker of hepatocellular differentiation in fine needle aspiration cytology and a marker of hepatocytes and hepatocellular neoplasms. ARG1 is closely associated with alternative macrophage activation(PMID:12098359) and ARG1 has been shown to protect motor neurons from trophic factor deprivation and allow sensory neurons to overcome neurite outgrowth inhibition by myelin proteins(PMID: 20071539). It can exist as a homotrimer(PMID:16141327) and it has 3 isoforms produced by alternative splicing. Defects in ARG1 are the cause of argininemia (ARGIN). Deletion or TNF-mediated restriction of ARG1 unleashes the production of NO by NOS2, which is critical for pathogen control. (PMID:27117406). ARG1 is a novel biomarker of post-stroke immune suppression(2016). The antibody is conjugated with CL594, Ex/Em 593 nm/614 nm.

## Publications notables

Autrice	Pubmed ID	Journal	Application
Ying Kong	34258169	Adv Sci (Weinh)	IF

## Stockage

Stockage:  
Stocker à -20 °C. Éviter toute exposition à la lumière. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.  
L'aliquotage n'est pas nécessaire pour le stockage à -20°C

\*\*\* Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:  
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

---

## Données de validation sélectionnées



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CL594-66129 (ARG1 antibody) at dilution of 1:50.