

À des fins de recherche uniquement

Anticorps Monoclonal anti-DEK

Numéro de catalogue: **CL594-66194** **Phare**



Informations de base

Numéro de catalogue: CL594-66194	Numéro d'acquisition GenBank: BC035259	Méthode de purification: Purification par protéine A
Taille: 100ul , Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI): 7913	CloneNo.: 5E3A6
Hôte: Mouse	Nom complet: DEK oncogene	Dilutions recommandées: WB 1:500-1:1000
Isotype: IgG2a	MW calculé: 43 kDa	Excitation/Emission maxima wavelengths: 588 nm / 604 nm
Immunogen Catalog Number: AG8225	MW observés: 50 kDa	

Applications

Applications testées: FC (Intra), WB	Contrôles positifs: WB : cellules HEK-293, cellules HepG2, cellules Jurkat
Spécificité de l'espèce: Humain	

Informations générales

Dek oncogene(DEK) is first recognized in a fusion with CAN nucleoporin protein, and it's a also autoantigen in patients with pquiarticular onset juvenile rheumatoid arthritis. As a site-specific binding protein, DEK may involve in transcriptional regulation and signal transduction. Analysis by agarose gel electrophoresis, DEK showed to reduce the efficiency with which chromatin is replicated. It's one of the first components of the splicing complex, which remains associate with spliced exons dependent on prior splicing of pre-mRNA.

Stockage

Stockage:
Stocker à -20 °C. Éviter toute exposition à la lumière. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.
L'aliquotage n'est pas nécessaire pour le stockage à -20C

***** Les 20ul contiennent 0,1% de BSA.**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with CL594-66194 (DEK antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



1×10^6 HeLa cells were intracellularly stained with 0.4 μ g CoraLite[®]594 Anti-Human DEK (CL594-66194, Clone:5E3A6) (red), or 0.4 μ g Mouse IgG2a Isotype Control (CL594-66360-2, Clone: K11A1B2A2) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).