

À des fins de recherche uniquement

Anticorps Monoclonal anti-STAT5B

Numéro de catalogue:**CL594-66427**



Informations de base

| | | |
|--|---|---|
| Numéro de catalogue: | BC020868 | Méthode de purification: |
| CL594-66427 | Purification par protéine A | |
| Taille: | Identification du gène (NCBI): | CloneNo.: |
| 100ul , Concentration: 1000 µg/ml by Nanodrop; | 6777 | 1D9B11 |
| Hôte: | Nom complet: | Dilutions recommandées: |
| Mouse | signal transducer and activator of transcription 5B | IF 1:50-1:500 |
| Isotype: | MW calculé | Excitation/Emission maxima wavelengths: |
| IgG2a | 787 aa, 90 kDa | 588 nm / 604 nm |
| Immunogen Catalog Number: | MW observés: | |
| AG2709 | 90 kDa | |

Applications

| | |
|--------------------------|----------------------|
| Applications testées: | Contrôles positifs: |
| FC (Intra), IF | IF : cellules HepG2, |
| Spécificité de l'espèce: | |
| Humain | |

Informations générales

STAT5B belongs to the transcription factor STAT family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. STAT5B carries out a dual function: signal transduction and activation of transcription. STAT5B mediates the signal transduction triggered by various cell ligands, such as IL2, IL4, CSF1, and different GH. It has been shown to be involved in diverse biological processes, such as TCR signalling, apoptosis, adult mammary gland development, and sexual dimorphism of liver gene expression. Despite the 95% homology between the immunogen (Ag2709) and the STAT5A protein, our WB detection showed that this antibody does not recognize the STAT5A fusion protein (Ag19546).

Stockage


Stockage:
Stocker à -20 °C. Éviter toute exposition à la lumière. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.
L'aliquotage n'est pas nécessaire pour le stockage à -20°C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Coralite@594 STAT5B antibody (CL594-66427, Clone: 1D9B11) at dilution of 1:200, CL488-Phalloidin (green).



1X10⁶ HepG2 cells were intracellularly stained with 0.4 ug Coralite@594 Anti-Human STAT5B (CL594-66427, Clone:1D9B11) (red), or 0.4 ug Mouse IgG2a Isotype Control (CL594-66360-2, Clone: K11A1B2A2) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).