

À des fins de recherche uniquement

Anticorps Monoclonal anti-FUCA1

Numéro de catalogue:**CL594-67251**



Informations de base

| | | |
|--|--------------------------------|----------------------------|
| Numéro de catalogue: | BC017338 | Méthode de purification: |
| CL594-67251 | Purification par protéine A | |
| Taille: | Identification du gène (NCBI): | CloneNo.: |
| 100ul , Concentration: 1000 µg/ml by Nanodrop; | 2517 | 1G5C5 |
| Hôte: | Nom complet: | Dilutions recommandées: |
| Mouse | fucosidase, alpha-L- 1, tissue | IF 1:50-1:500 |
| Isotype: | MW calculé | Excitation/Emission maxima |
| IgG2b | 461 aa, 53 kDa | wavelengths: |
| Immunogen Catalog Number: | MW observés: | 588 nm / 604 nm |
| AG9638 | 50-56 kDa | |

Applications

| | |
|--------------------------|--------------------------------------|
| Applications testées: | Contrôles positifs: |
| IF | IF : tissu de cancer du foie humain, |
| Spécificité de l'espèce: | Humain, souris |

Informations générales

FUCA1(a-L-Fucosidase) is a lysosomal glycosidase which is ubiquitous in eukaryotic cells. This enzyme is usually found as a soluble component of the lysosome and functions as an acid hydrolase in the degradation of numerous and diverse fucoglycoconjugates. The mammalian a-L-fucosidases are relatively large, multisubunit glycoproteins which exist in multiple molecular isoforms. This enzyme may have a role in sperm-egg interactions in the reproductive tract of the female rat. There is a precursor 54 kDa form of rat sperm fucosidase which is processed to a mature 52 kDa form of the enzyme (PMID:8836125). This protein from rat epididymis is a tetramer of 210-220 kDa, made up by two pairs of subunits of 47 and 60 kDa, and the one from human seminal plasma exists in different oligomeric forms, that is, as a dimer, as a tetramer, and as a hexamer made up of a single subunit of 56-57 kDa (PMID:16736526).

Stockage


Stockage:
Stocker à -20 °C. Éviter toute exposition à la lumière.
Tampon de stockage:
PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.
L'aliquotage n'est pas nécessaire pour le stockage à -20°C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Immunofluorescent analysis of (4% PFA) fixed human liver cancer tissue using Coralite®594 FUCA1 antibody (CL594-67251, Clone: 1G5C5) at dilution of 1:200. DAPI (blue).



Immunofluorescent analysis of (4% PFA) fixed human liver cancer tissue using Coralite®594 FUCA1 antibody (CL594-67251, Clone: 1G5C5) at dilution of 1:200. DAPI (blue).