

À des fins de recherche uniquement

Anticorps Monoclonal anti-ACC1

Numéro de catalogue: **CL594-67373**



Informations de base

| | | |
|--|--|---|
| Numéro de catalogue: CL594-67373 | Numéro d'acquisition GenBank: BC137287 | Méthode de purification: Purification par protéine A |
| Taille: 100ul , Concentration: 1000 µg/ml by Nanodrop; | Identification du gène (NCBI): 31 | CloneNo.: 1A11G10 |
| Hôte: Mouse | Nom complet: acetyl-Coenzyme A carboxylase alpha | Dilutions recommandées: IF 1:50-1:500 |
| Isotype: IgG2a | MW calculé 2383 aa, 275 kDa | Excitation/Emission maxima wavelengths: 588 nm / 604 nm |
| Immunogen Catalog Number: AG17503 | MW observés: 260-270 kDa | |

Applications

| | |
|---|---|
| Applications testées: FC (Intra), IF | Contrôles positifs: IF : cellules HepG2, |
| Spécificité de l'espèce: Humain | |

Informations générales

ACACA (Acetyl-CoA carboxylase 1, ACC), also named as ACAC, ACC1 and ACCA, belongs to the biotin containing enzyme family. It catalyzes the synthesis of malonyl-CoA, which is an intermediate substrate playing a pivotal role in the regulation of fatty acid metabolism and energy production. ACACA is involved in the biosynthesis of fatty acids, and malonyl-CoA produced is used as a building block to extend the chain length of fatty acids by fatty acid synthase (FAS) (PMID:19900410).

Stockage

Stockage:
Stocker à -20 °C. Éviter toute exposition à la lumière.
Tampon de stockage:
PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.
L'aliquotage n'est pas nécessaire pour le stockage à -20C

***** Les 20ul contiennent 0,1% de BSA.**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CoraLite®594-conjugated ACC1 antibody (CL594-67373, Clone: 1A11G10) at dilution of 1:100.



1X10⁶ HeLa cells were intracellularly stained with 0.4 ug CoraLite®594 Anti-Human ACC1 (CL594-67373, Clone:1A11G10) (red), or 0.4 ug Mouse IgG2a Isotype Control (CL594-66360-2, Clone: K11A1B2A2) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).