

À des fins de recherche uniquement

# Anticorps Polyclonal de lapin anti-NONO



Numéro de catalogue: **CL647-11058**

Phare

## Informations de base

<b>Numéro de catalogue:</b> CL647-11058	<b>Numéro d'acquisition GenBank:</b> BC010049	<b>Méthode de purification:</b> Purification par affinité contre l'antigène
<b>Taille:</b> 100ul , Concentration: 1000 ug/mL by Nanodrop;	<b>Identification du gène (NCBI):</b> 4841	<b>Excitation/Emission maxima wavelengths:</b> 654 nm / 674 nm
<b>Hôte:</b> Lapin	<b>Nom complet:</b> non-POU domain containing, octamer-binding	
<b>Isotype:</b> IgG	<b>MW calculé:</b> 54 kDa	
<b>Immunogen Catalog Number:</b> AG1512	<b>MW observés:</b> 54 kDa	

## Applications

**Applications testées:**  
FC (Intra)

**Spécificité de l'espèce:**  
Humain, rat, souris

## Informations générales

Non-pou domain-containing octamer-binding protein(NONO), other named p54nrb, belongs to DBHS family, whose protein are predominately nuclear and involves in transcriptional and posttranscriptional gene regulatory functions as well as DNA repair. This family has roles in a wide range of biological processes, such as the regulation of circadian rhythm, carcinogenesis and progression of cancer. NONO forms heterodimer with SFPO, and has been implicated in nuclear retention of A- to I-edited RNA as paraspeckle components, pre-mRAN 3'-end formation, cAMP cycling and transcriptional activation.

## Stockage

**Stockage:**  
Stocker à -20 °C. Éviter toute exposition à la lumière. Stable pendant un an après l'expédition.

**Tampon de stockage:**  
PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.  
L'aliquotage n'est pas nécessaire pour le stockage à -20C

**\*\*\* Les 20ul contiennent 0,1% de BSA.**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Données de validation sélectionnées



1X10<sup>6</sup> HepG2 cells were intracellularly stained with 0.2 ug CoraLite® Plus 647 Anti-Human NONO (CL647-11058) (red), or 0.2 ug isotype control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).