

À des fins de recherche uniquement

# Anticorps Polyclonal de lapin anti- EIF2S1/EIF2A

Numéro de catalogue:[CL647-11170](#)



## Informations de base

Numéro de catalogue:	BC002513	Méthode de purification:
CL647-11170		Purification par affinité contre l'antigène
Taille:	Identification du gène (NCBI):	Excitation/Emission maxima wavelengths:
100µl , Concentration: 1000 µg/ml by Nanodrop;	1965	654 nm / 674 nm
Hôte:	Nom complet:	
Lapin	eukaryotic translation initiation factor 2, subunit 1 alpha, 35kDa	
Isotype:	MW calculé	
IgG	36 kDa	
Immunogen Catalog Number:	MW observés:	
AG1645	36 kDa	

## Applications

### Applications testées:

FC (Intra)

### Spécificité de l'espèce:

Humain, rat, souris

## Informations générales

EIF2S1 is one subunit of the translation initiation factor EIF2, which catalyzes the first regulated step of protein synthesis initiation, promoting the binding of the initiator tRNA to 40S ribosomal subunits. This complex binds to 40S ribosomal subunit, followed by mRNA binding to form a 43S preinitiation complex. Junction of the 60S ribosomal subunit to form the 80S initiation complex is preceded by hydrolysis of the GTP bound to eIF-2 and release of an eIF-2-GDP binary complex. In order for eIF-2 to recycle and catalyze another round of initiation, the GDP bound to eIF-2 must exchange with GTP by way of a reaction catalyzed by eIF-2B. EIF2A (Gene ID: 83939) and EIF2S1 (Gene ID: 1965) share the EIF2A symbol/alias in common. EIF2S1 is the alpha subunit of the EIF2 translation initiation complex. Although both of these proteins function in binding initiator tRNA to the 40S ribosomal subunit, the EIF2A protein does so in a codon-dependent manner, whereas EIF2 complex requires GTP.

## Stockage

### Stockage:

Stocker à -20 °C. Éviter toute exposition à la lumière. Stable pendant un an après l'expédition.

### Tampon de stockage:

PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.


L'aliquotage n'est pas nécessaire pour le stockage à -20°C

\*\*\* Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:  
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)      E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Données de validation sélectionnées



1X10<sup>6</sup> HeLa cells were intracellularly stained with 0.2 ug Coralite® Plus 647 Anti-Human EIF2S1/EIF2A (CL647-11170) (red), or 0.2 ug CL647-30000 Rabbit IgG (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).