

À des fins de recherche uniquement

Anticorps Monoclonal anti-MHC Class II (I-A/I-E)



Numéro de catalogue:**CL647-65122** 1 Publications

Informations de base

Numéro de catalogue:	BC031711	Méthode de purification:
CL647-65122	Purification par affinité	
Taille:	14960	CloneNo.:
100ug , 0.5 mg/ml	M5/114.15.2	
Hôte:	histocompatibility 2, class II antigen	Dilutions recommandées:
Rat	A, alpha	IF 1:200-1:800
Isotype:		Excitation/Emission maxima wavelengths:
IgG2b, kappa		654 nm / 674 nm

Applications

Applications testées:	Contrôles positifs:
FC, IF	IF : splénocytes de souris,
Demandes citées:	
FC	
Spécificité de l'espèce:	
souris	
Espèces citées:	
souris	

Informations générales

Major histocompatibility complex (MHC) class II molecules are heterodimeric transmembrane glycoproteins that are expressed on the surface of antigen-presenting cells such as dendritic cells, macrophages, and B cells. The M5/114.15.2 antibody detects polymorphic determinants present on B cells, monocytes, macrophages, dendritic cells, and activated T lymphocytes from mice carrying the H-2b, H-2d, H-2q, H-2p, H-2r and H-2u haplotypes but not from mice carrying the H-2s or H-2f haplotypes (PMID: 6170707). This antibody recognizes both I-A and I-E subregion-encoded Ia glycoproteins (I-Ab, I-Ad, I-Aq, I-Ed, I-Ek, not I-Af, I-Ak, or I-As). M5/114 mAb is reported to inhibit I-A subregion-restricted T cell responses of the H-2b, H-2d, H-2q, H-2u but not the H-2f, H-2k, H-2s haplotypes (PMID: 6173436).

Publications notables

Autrice	Pubmed ID	Journal	Application
Luqiu Feng	36918933	Vet Res	FC


Stockage

Stockage:
Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.
Tampon de stockage:
PBS avec azoture de sodium à 0,1 % et BSA à 0,5 %, pH 7,3.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1X10⁶ BALB/c mouse splenocytes were surface stained with CoraLite®488 Anti-Mouse CD3 (CL488-65077, Clone: 17A2) and 0.25 ug CoraLite® Plus 647 Anti-Mouse MHC Class II (I-A/I-E) (CL647-65122, Clone: M5/114.15.2) or CoraLite® Plus 647-Conjugated rat IgG2b isotype control. Cells were not fixed.



Immunofluorescent analysis of mouse splenocytes using CoraLite® Plus 647-conjugated Anti-Mouse MHC Class II (I-A/I-E) (CL647-65122, Clone: M5/114.15.2) at dilution of 1:400.