## À des fins de recherche uniquement

## Anticorps Monoclonal anti-CD64

Numéro de catalogue: CL647-65253



Informations de base

Numéro de catalogue:

CL647-65253

Taille:

100 tests , 5  $\mu$ l/test

Hôte:

Mouse Isotype:

IgG1, kappa

Numéro d'acquisition GenBank:

BC032634

Identification du gène (NCBI):

2209

Nom complet: Fc fragment of IgG, high affinity Ia,

receptor (CD64)

MW calculé 374 aa, 43 kDa Méthode de purification: Purification par affinité

CloneNo.:

Excitation/Emission maxima

wavelengths: 654 nm / 674 nm

**Applications** 

Applications testées:

Spécificité de l'espèce:

Humain

Informations générales

Fcy receptor comprise a multigene family of integral membrane glycoproteins that exhibit complex activation or  $inhibitory\ effects\ on\ cell\ functions\ after\ aggregation\ by\ complexed\ immunoglobulin\ G\ (IgG)\ (PMID:\ 17005690\ ).$ CD64, also known as Fcy RIA, is a high-affinity receptor for the Fc region of IgG. It is expressed by monocytes/macrophages, activated neutrophils, dendritic cells, and early myeloid cells (PMID: 23293080; 19642859; 7680917). CD64 functions in both innate and adaptive immune responses.

Stockage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Tampon de stockage:

PBS with 0.09% sodium azide and 0.5% BSA.


in USA), or 1(312) 455-8498 (outside USA)

W: ptglab.com

## Données de validation sélectionnées



1X10^6 human PBMCs were surface stained with 5 ul Coralite® Plus 647 Anti-Human CD64 (CL647-65253, Clone:10.1) or unstained. Cells were not fixed. Monocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul Coralite® Plus 647 Anti-Human CD64 (Cl.647-65253, Clone:10.1) or unstained. Cells were not fixed. Monocytes were gated.