

À des fins de recherche uniquement

Anticorps Monoclonal anti-M-CSF

Numéro de catalogue: **CL647-65283**



Informations de base

Numéro de catalogue:

CL647-65283

Taille:

100ug, 0.2 mg/mL

Hôte:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

HZ-1192

Numéro d'acquisition GenBank:

BC021117

Identification du gène (NCBI):

1435

Nom complet:

colony stimulating factor 1
(macrophage)

MW calculé

60 kDa

Méthode de purification:

Purification par affinité

CloneNo.:

3G12E6

Excitation/Emission maxima
wavelengths:

654 nm / 674 nm

Applications

Applications testées:

FC

Spécificité de l'espèce:

Humain

Informations générales

M-CSF plays an essential role in the regulation of survival, proliferation and differentiation of hematopoietic precursor cells, especially mononuclear phagocytes. M-CSF promotes the release of proinflammatory chemokines, and thereby plays an important role in innate immunity and in inflammatory processes. It also plays an important role in the regulation of osteoclast proliferation and differentiation, the regulation of bone resorption, and is required for normal bone development. On the cellular level, M-CSF promotes reorganization of the actin cytoskeleton, regulates formation of membrane ruffles, cell adhesion and cell migration and plays a role in lipoprotein clearance (PMID: 29734839; 29679908; 28202039)

Stockage

Stockage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Tampon de stockage:

PBS with 0.09% sodium azide and 0.5% BSA.



For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1X10⁶ LPS and TNF- α then Brefeldin A treated human PBMCs were surface stained with CoraLite® Plus 555 Anti-Human CD11b then intracellularly stained with 0.2 ug CoraLite® Plus 647 Anti-Human M-CSF (CL647-65283, Clone:3G12E6) or Mouse IgG1 Isotype Control. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). Monocytes were gated.

1X10⁶ LPS and TNF- α then Brefeldin A treated human PBMCs were intracellularly stained with 0.2 ug CoraLite® Plus 647 Anti-Human M-CSF (CL647-65283, Clone:3G12E6) (red) or Mouse IgG1 Isotype Control (blue), or unactivated then Brefeldin A treated human PBMCs were intracellularly stained with 0.2 ug CoraLite® Plus 647 Anti-Human M-CSF (CL647-65283, Clone:3G12E6) (black, dashed line). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit

