

À des fins de recherche uniquement

# Anticorps Monoclonal anti-Alpha Tubulin

Numéro de catalogue:**CL647-66031**



## Informations de base

Numéro de catalogue:	BC009314	Méthode de purification:
CL647-66031	Purification par protéine A	
Taille:	Identification du gène (NCBI):	CloneNo.:
100ul , Concentration: 1000 µg/ml by Nanodrop;	10376	1E4C11
Hôte:	Nom complet:	Dilutions recommandées:
Mouse	tubulin, alpha 1b	IF 1:50-1:500
Isotype:	MW calculé	Excitation/Emission maxima
IgG2b	50 kDa	wavelengths:
Immunogen Catalog Number:	MW observés:	654 nm / 674 nm
AG18034	50-55 kDa	

## Applications

Applications testées:	Contrôles positifs:
FC (Intra), IF	IF : cellules HeLa,
Spécificité de l'espèce:	
canin, Humain, rat, souris	

## Informations générales

There are five tubulins in human cells: alpha, beta, gamma, delta, and epsilon. Tubulins are conserved across species. They form heterodimers, which multimerize to form a microtubule filament. An alpha and beta tubulin heterodimer is the basic structural unit of microtubules. The heterodimer does not come apart once formed. The alpha and beta tubulins, which are each about 55 kDa MW, are homologous but not identical. Alpha tubulin is useful for scientists across fields as an internal control due to its high, ubiquitous expression pattern. Tubulin expression may vary according to resistance to antimicrobial and antimitotic drugs. This antibody specifically recognizes Tubulin Alpha.

## Stockage


**Stockage:**  
Stocker à -20 °C. Éviter toute exposition à la lumière. Stable pendant un an après l'expédition.  
**Tampon de stockage:**  
PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.  
**L'aliquotage n'est pas nécessaire pour le stockage à -20°C**

\*\*\* Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:  
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)      E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Données de validation sélectionnées



1X10<sup>6</sup> HepG2 cells were intracellularly stained with 0.4 ug Coralite® Plus 647 Anti-Human Alpha Tubulin (CL647-66031, Clone:1E4C11) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).

Immunofluorescent analysis of (4% PFA) fixed HeLa cells using Coralite® Plus 647 Alpha Tubulin antibody (CL647-66031, Clone: 1E4C11 ) at dilution of 1:200.