

À des fins de recherche uniquement

Anticorps Monoclonal anti-GNAS

Numéro de catalogue: CL647-66253



Informations de base

Numéro de catalogue: CL647-66253	Numéro d'acquisition GenBank: BC002722	Méthode de purification: Purification par protéine A
Taille: 100ul, Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI): 2778	CloneNo.: 1A9C10
Hôte: Mouse	Nom complet: GNAS complex locus	Excitation/Emission maxima wavelengths: 654 nm / 674 nm
Isotype: IgG2b	MW calculé 45 kDa	
Immunogen Catalog Number: AG20530	MW observés: 46 kDa	

Applications

Applications testées:
FC (Intra)
Spécificité de l'espèce:
Humain, porc

Informations générales

Guanine nucleotide binding protein (G protein), alpha stimulating activity polypeptide 1 (GNAS1) is the ubiquitously expressed heterotrimeric G protein that couples receptors to the effector enzyme adenylyl cyclase and is required for receptor-stimulated intracellular cAMP generation. Mutations of Gs(alpha) residues involved in the GTPase reaction that lead to constitutive activation are present in endocrine tumors, fibrous dysplasia of bone, and McCune-Albright syndrome. The molecular weight of Gs(alpha) protein is about 46 kDa.

Stockage

Stockage:
Stocker à -20 °C. Éviter toute exposition à la lumière. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.
L'aliquotage n'est pas nécessaire pour le stockage à -20C

***** Les 20ul contiennent 0,1% de BSA.**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1X10⁶ MCF-7 cells were intracellularly stained with 0.2 ug CoraLite® Plus 647 Anti-Human GNAS (CL647-66253, Clone:1A9C10) (red), or 0.2 ug isotype control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).