

À des fins de recherche uniquement

# Anticorps Monoclonal anti-NQO1

Numéro de catalogue:**CL647-67240**



## Informations de base

Numéro de catalogue:	BC007659	Méthode de purification:
CL647-67240	Purification par protéine A	
Taille:	Identification du gène (NCBI):	CloneNo.:
100ul , Concentration: 1000 µg/ml by Nanodrop;	1728	1E5G7
Hôte:	Nom complet:	Excitation/Emission maxima
Mouse	NAD(P)H dehydrogenase, quinone 1	wavelengths:
Isotype:	MW calculé	654 nm / 674 nm
IgG2a	274 aa, 31 kDa	
Immunogen Catalog Number:	MW observés:	
AG28933	29-31 kDa	

## Applications

### Applications testées:

FC (Intra)

### Spécificité de l'espèce:

Humain, porc, rat

## Informations générales

NQO1, also named as DIA4, NMOR1, DTD and QR1, belongs to the NAD(P)H dehydrogenase (quinone) family. This enzyme apparently serves as a quinone reductase in connection with conjugation reactions of hydroquinones involved in detoxification pathways as well as in biosynthetic processes such as the vitamin K-dependent gamma-carboxylation of glutamate residues in prothrombin synthesis. It is known to be involved in benzene metabolism. In human studies of ozone exposure, polymorphisms in oxidative stress genes (NQO1, GSTM1, GSTP1) modify respiratory symptoms, lung function, biomarkers and risk of asthma. (PMID:18511640; 18848868 )

## Stockage

### Stockage:

Stocker à -20 °C. Éviter toute exposition à la lumière. Stable pendant un an après l'expédition.

### Tampon de stockage:

PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.


L'aliquotage n'est pas nécessaire pour le stockage à -20°C

\*\*\* Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:  
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)      E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Données de validation sélectionnées



1X10<sup>6</sup> MCF-7 cells were intracellularly stained with 0.2 ug CoralLite® Plus 647 Anti-Human NQO1 (CL647-67240, Clone:1E5G7) (red), or 0.2 ug isotype control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).