

À des fins de recherche uniquement

Anticorps Monoclonal anti-MFN2

Numéro de catalogue:**CL647-67487**

Phare



Informations de base

Numéro de catalogue:	BC017061	Méthode de purification:
CL647-67487	Purification par protéine A	
Taille:	Identification du gène (NCBI):	CloneNo.:
100ul , Concentration: 1000 µg/ml by Nanodrop;	9927	5F3B3
Hôte:	Nom complet:	Excitation/Emission maxima
Mouse	mitofusin 2	wavelengths:
Isotype:	MW calculé	654 nm / 674 nm
IgG2a	757 aa, 86 kDa	
Immunogen Catalog Number:	MW observés:	
AG29873	86 kDa	

Applications

Applications testées:

FC (Intra)

Spécificité de l'espèce:

Humain, rat, souris

Informations générales

MFN2, also named as CPRP1 and KIAA0214, belongs to the mitofusin family. It is an Essential transmembrane GTPase, which mediates mitochondrial fusion. MFN2 acts independently of the cytoskeleton. It therefore plays a central role in mitochondrial metabolism and may be associated with obesity and/or apoptosis processes. Overexpression of MFN2 induces the formation of mitochondrial networks. It plays an important role in the regulation of vascular smooth muscle cell proliferation. Defects in MFN2 are the cause of Charcot-Marie-Tooth disease type 2A2 (CMT2A2). Defects in MFN2 are the cause of Charcot-Marie-Tooth disease type 6 (CMT6). Ubiquitinated forms of Mfn2 (mono- and polyubiquitinated) are present during mitophagy.

Stockage

Stockage:

Stocker à -20 °C. Éviter toute exposition à la lumière.

Tampon de stockage:

PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.


L'aliquotage n'est pas nécessaire pour le stockage à -20°C

*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1X10⁶ HepG2 cells were intracellularly stained with 0.2 ug CoralLite® Plus 647 Anti-Human MFN2 (CL647-67487, Clone:5F3B3) (red), or 0.2 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).