À des fins de recherche uniquement

Anticorps Monoclonal anti-CD73

Numéro de catalogue:CL750-65162



Informations de base

Numéro de catalogue:

CL750-65162

Taille:

100 tests , 5 μ l/test

Hôte:

Mouse Isotype:

Mouse IgG1 kappa

Applications testées:

FC

Spécificité de l'espèce:

Humain

Numéro d'acquisition GenBank:

Méthode de purification: Purification par protéine G

CloneNo.:

AD2

Excitation/Emission maxima

wavelengths: 755 nm / 780 nm

Informations générales

Applications

CD73, also known as ecto-5'-nucleotidase (5'-NT), is a 70-kDa, glycosyl-phosphatidylinositol-linked membranebound glycoprotein found in most tissues (PMID: 18404475; 20179192). CD73 is an ectoenzyme that catalyzes the dephosphorylation of AMP and other nucleoside monophosphates (PMID: 9553767). In the human immune system, CD73 is expressed on subsets of T and B cells, on germinal center follicular dendritic cells, and on thymic medullary reticular fibroblasts and epithelial cells (PMID: 2137649; 9553767). CD73 is highly expressed in many human solid tumors and is closely involved in cancer progression (PMID: 20179192).

Stockage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

BC015940

Nom complet:

MW calculé

29 kDa, 63 kDa


Identification du gène (NCBI):

5'-nucleotidase, ecto (CD73)


Tampon de stockage:

PBS with 0.09% sodium azide and 0.5% BSA.

Données de validation sélectionnées



1X10^6 human PBMCs were surface co-stained with FITC Anti-Human CD3 and 5 ul Coralite® Plus 750 Anti-Human CD73 (CL750-65162, Clone:AD2) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul Coralite® Plus 750 Anti-Human CD73 (CL750-65162, Clone:AD2) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.