

À des fins de recherche uniquement

Anticorps Monoclonal anti-CD64

Numéro de catalogue:**CL750-65253**



Informations de base

Numéro de catalogue:	BC032634	Méthode de purification:
CL750-65253	Purification par affinité	
Taille:	Identification du gène (NCBI):	CloneNo.:
100tests , 5 µl/test	2209	10.1
Hôte:	Nom complet:	Excitation/Emission maxima
Mouse	Fc fragment of IgG, high affinity Ia, receptor (CD64)	wavelengths:
Isotype:	MW calculé	755 nm / 780 nm
IgG1, kappa	374 aa, 43 kDa	

Applications

Applications testées:
FC

Spécificité de l'espèce:
Humain

Informations générales

Fcγ receptor comprise a multigene family of integral membrane glycoproteins that exhibit complex activation or inhibitory effects on cell functions after aggregation by complexed immunoglobulin G (IgG) (PMID: 17005690). CD64, also known as Fcγ RIA, is a high-affinity receptor for the Fc region of IgG. It is expressed by monocytes/macrophages, activated neutrophils, dendritic cells, and early myeloid cells (PMID: 23293080; 19642859; 7680917). CD64 functions in both innate and adaptive immune responses.


Stockage

Stockage:
Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.
Tampon de stockage:
PBS with 0.09% sodium azide and 0.5% BSA.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1×10^6 human PBMCs were surface stained with 5 μ l Coralite® Plus 750 Anti-Human CD64 (CL750-65253, Clone:10.1). Cells were not fixed. Monocytes were gated.