

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-Cyclin E2

Numéro de catalogue:**CR-11935**



Informations de base

Numéro de catalogue:	BC020729	Méthode de purification:
CR-11935		Purification par affinité contre l'antigène
Taille:	Identification du gène (NCBI):	Excitation/Emission maxima wavelengths:
100ul , Concentration: 1000 µg/ml by Nanodrop;	9134	592 nm / 611 nm
Hôte:	Nom complet:	
Lapin	cyclin E2	
Isotype:	MW calculé	
IgG	374 aa, 44 kDa	
Immunogen Catalog Number:	MW observés:	
AG2532	44 kDa	

Applications

Applications testées:

FC (Intra)

Spécificité de l'espèce:

Humain

Informations générales

Cyclin E2 (CCNE2) belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance throughout the cell cycle. Cyclins function as regulators of cyclin-dependent kinases (CDKs). Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of cell cycle events. CCNE2 forms a complex with and functions as a regulatory subunit of CDK2 and has been shown to specifically interact with CIP/KIP family of CDK inhibitors. CCNE2 plays a role in cell cycle G1/S transition and its expression peaks at the G1-S phase. Whereas cyclin E1 is expressed in most proliferating normal and tumor cells, cyclin E2 levels are low or undetectable in nontransformed cells, and are elevated in tumor-derived cells.

Stockage

Stockage:

Stocker à -20 °C. Éviter toute exposition à la lumière. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.


L'aliquotage n'est pas nécessaire pour le stockage à -20°C

*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1X10⁶ HeLa cells were intracellularly stained with 0.2 ug Cardinal Red™ Anti-Human Cyclin E2 (CR-11935) (red), or 0.2 ug isotype control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).